

**Review of Maternal Deprivation Experiments on Primates
at the National Institutes of Health**

Prepared by

Katherine Roe, Ph.D.

Research Associate

Laboratory Investigations Department

People for the Ethical Treatment of Animals

This document provides a critical scientific review and assessment of continuing maternal deprivation and psychopathology studies on nonhuman primates conducted within the National Institutes of Health (NIH) Intramural Research Program. A careful analysis of Animal Study Proposals, Board of Scientific Counselors reviews, scientific publications, photographs, and videos related to these projects casts doubt on the worth of these experiments in light of advancements in the field, and offers several examples of human-based studies that successfully address precisely the questions asked by these NIH investigators. Moreover, after consulting numerous experts in the fields of anthropology, primatology, medicine, and mental health, we conclude **that given the harm caused to animals, the experiments' limited relevance to humans, the substantial financial cost, and the existence of superior nonanimal research methods that the continued use of animals in this work is scientifically and ethically unjustifiable.**

Project title: “Biobehavioral Reactivity in Monkeys”

Institute: National Institute of Child Health and Development (NICHD)

Principal Investigator: Stephen J. Suomi

Intramural Animal Study Proposals: 11-043, 14-043

Project Number: 1ZIAHD001106

Start/end: 2007–present

Funding: \$907,723 in 2013 (\$7,786,372 total)

At the foundation of all of the studies in question are maternal deprivation experiments conducted by Stephen J. Suomi and the Laboratory of Cognitive Ethology (LCE) at NICHD. For the past three decades, Suomi’s group has utilized a maternal deprivation model of psychopathology, depriving hundreds of infant macaques of maternal contact and resulting in animals with an array of cognitive, social, emotional, and physical deficits that persist throughout their lifetimes. According to the approved Animal Study Proposal (ASP), approximately 45 macaques are selectively bred each year to carry different alleles of the 5H-TTT and MAO-1 genes, known to be risk factors for psychopathology in humans. Half of these captive-born infants are separated from their mothers within 24 hours of birth, causing great distress to mother and baby, and are hand-reared by humans in a nursery for one month and then put into a nursery with other like-reared peers, sometimes with a terrycloth-covered water bottle. Starting on their first day of birth, all infants are subject to numerous fear, stress, and pain-inducing tests. Day-old infants are forcibly restrained by experimenters for behavioral tests, such as facial imitation or head-orientation bias trials. Other experiments entail the infants being isolated in small cages, placed in unfamiliar locations, and deliberately startled by threatening human strangers, unfamiliar objects (including realistic-looking snakes, which are innately frightening to monkeys), and unfamiliar conspecifics. In one such procedure designed to measure infants’ auditory startle response, newborn infants are restrained inside tiny mesh cages and placed in “startle chambers” where they are presented with unexpected loud noises. During their first few months of life, the infants are repeatedly subjected to blood draws and cerebral spinal fluid taps; hair and saliva samples are also taken. Additionally, in a project funded by the NICHD (Project 5P01HD064653; \$877,229 of funding in 2013), Nathan A. Fox from the University of Maryland takes infants as young as one day old from Suomi’s colony, shaves their heads, and physically restrains them for electroencephalogram testing.^{1,2}

The approved ASPs for the breeding and experimentation regimen (11-043, 14-043) in Suomi’s laboratory does not explain the scientific relevance of the single nucleotide polymorphisms that animals are bred to carry, their methods for selective breeding of these animals, the exact conditions they classify as “mother-rearing,” the scientific purpose for numerous cognitive and biological tests

being conducted, or any risk factors associated with capture, restraint, and biological or behavioral testing that they perform repeatedly on the animals.

The NIH Policy Manual for Animal Care and Use in the Intramural Research Program clearly states that the Principal/Responsible Investigator is accountable for assuring that the “proposed studies are not unnecessarily duplicative” (p. 7).³ Several of the experiments currently being conducted have already been performed using the same procedures and the results published.^{4,5,6,7,8,9,10,11,12} The rearing procedures described have been in place for decades, and behavioral and biological data from these animals have also been collected for decades.^{13,14,15,16} Repeating these test batteries and causing suffering to additional infant monkeys is required by law to be justified; however, given the limited information contained in the ASP, it is virtually impossible for a review committee to adequately evaluate the project’s design or scientific merit. The LCE’s approved ASP emphasizes that the purpose of the study is to model the genetic and environmental contributions to abnormal human behavior and to develop interventions for at-risk individuals. However, a comprehensive review shows that none of the aforementioned studies have resulted in the development or modification of treatments for the human mental illnesses they are purported to model.

In addition to the study designed to create and quantify mental illness in infant macaques, the LCE has also received \$6,289,327 since 2007 to assess whether the laboratory-reared, mentally ill animals they created can adapt to a non-laboratory environment (Project 1ZIAHD001107). According to the approved ASPs associated with this project (11-105, 14-105), the purpose of the study is to understand “how humans of all ages and backgrounds adapt to new physical and social settings, as well as what aspects of their immediate environment might be affecting their psychological well-being.” However, in their 2013 annual NIH Intramural Database report, the experimenters describe several findings related to infant-mother communication, facial processing in infants, the effect of oxytocin on monkey-human interactions, and cortisol levels in nursing mothers’ milk. The discrepancy between the procedures and purposes outlined in the ASP and the reported findings from those procedures makes it difficult to evaluate the value of this study in understanding human health and behavior.

Though the ASPs for these projects claim the protocols are designed to elucidate genetic and environmental influences on pathological behavior unattainable with human participants, many resultant publications from these projects merely address whether macaques exhibit visual preferences, facial asymmetries, facial preferences, imitative behaviors, or similar hand- and head-orientation biases as those already well documented in human infants.^{17,18,19,20,21} Given the wealth of knowledge about human behaviors of this sort—and the non-invasive research with humans available to further explore these same issues—these studies are gratuitous.

Project title: “Assessment of Neural and Behavioral Alterations Associated with Chronic Fluoxetine Administration in Adolescence”

Institute: National Institute of Mental Health (NIMH)

Principal Investigator: Bruno Averbeck

Intramural Animal Study Proposal: IPC-01-09

Project Number: MH002902

Start/end: 2007–present

Funding: \$9,034,371 total

At NIMH, the Non-Human Primate Core purchases many of the maternally deprived, at-risk for illness animals created in Suomi's laboratory for its own battery of experiments. Some of these studies expose the animals to additional acute startle and isolation²² in hopes of eliciting a pathological response to stress as a function of their early-adverse rearing conditions. For example, infants and juveniles are restrained inside tiny mesh cages or in restraint chairs and placed into startle chambers where they are deliberately startled by the presence of a human, loud auditory stimuli, or powerful bursts of air. To acclimate them to the chair restraint, the older animals spend up to an hour a day, every day, strapped to a chair for weeks *prior* to testing. In other experiments, the infant monkeys are caged with their mothers—who are chemically sedated so as to be unresponsive—and placed in a car seat.²³ Videos of these experiments indicate that infants are terrified and confused while they try to revive their mothers.

In addition to various oral, subcutaneous, and intramuscular administrations of drugs, some animals are surgically implanted with devices that allow intracranial administration of pharmaceuticals, requiring multiple surgeries, weeks of recovery and pain management, and constant monitoring for infection. According to the ASP, the purpose of this pharmaceutical treatment is to “define specific neural pathways important to the expression of emotional, social, or cognitive deficits associated with differential rearing histories.” However, the exact drugs administered intracranially are not specified but described as “substances of interest [that] are likely to include NM concentrations of the neuropeptides oxytocin, vasopressin, CRH, MEK inhibitor PD98592, or GABA agonists such as muscimol and bicuculline, as well as genes attached to viral vectors (AAV-P11).” Without including this critical information in the ASP, there is no way for reviewers to evaluate the merits of the proposed experiments.

Some animals are injected with Interferon-alpha, which creates depressive-like symptoms in the monkeys and causes heightened sensitivity to pain, anhedonia, and anorexia. This procedure is classified as causing unrelieved pain and/or distress to those animals to whom it is administered. An unspecified number of animals in this project will be killed following pharmaceutical administration. In their approved ASP to conduct these experiments (IPC-01-09), the experimenters argue that “these experiments could provide important insights about the pathoetiology as well as potential, novel treatments for human syndromes with social detachment.” In their 2010 annual NIH Intramural Database report, they write, “A major public health concern has emerged regarding the treatment of children with psychotherapeutic drugs. This study seeks to inform this important concern.” However, these statements seem to contradict other claims from this same project in a subsequent publication in *The American Journal of Psychiatry* in which the authors themselves conclude the following: “...[M]any findings from behavioral and biochemical studies in monkeys and other animals are not replicated in humans. Accordingly, this study cannot directly address the safety and efficacy of SSRIs in children and adolescents with psychiatric disorders. ... [T]his animal model of maternal separation has never been validated as a measure of drug efficacy in humans[.] ... The only way to know definitively whether SSRIs persistently upregulate SERT in humans would be to study our species”(p. 7-8).²⁴

In addition to the projects and procedures described above, many animals from Suomi's LCE have been used for additional testing with the NIAAA. One project (Project Number: 1ZIAAA000214), which received \$4 million dollars between 2007 and 2010, studied juvenile monkeys' response to acute social separation,²⁵ spontaneous alcohol consumption,²⁶ and even acute ethanol exposure,²⁷ which requires the animals to be restrained while high concentrations of ethanol are administered

intravenously. These alcohol exposure studies often result in alcohol addiction, increased aggression, and increased susceptibility to depression in macaques.^{28,29,30,31} Other animals are transported to Wake Forest University to be used in Project 5U01AA014106 where they undergo additional alcohol exposure testing before being killed and dissected.^{32,33,34} The Wake Forest study received \$3,931,858 in funding from 2003 and 2011.

Inapplicability to human mental illness

The experimenters that are discussed above seek to justify the use of animals by positing that maternally deprived macaques model the effect of early-life stress on the development of mood and anxiety disorders in humans. In addition to fundamental differences in gene expression,^{35,36,37,38} brain anatomy and physiology,^{39,40,41,42} and development^{43,44} among humans and other primates, these adverse environments *do not* adequately represent the type of early social and physical stressors that precipitate mental illness in human children and adults. In reality, sexual abuse, physical abuse, prenatal stress, parental drug abuse, parental mental illness and/or criminal behavior, and economic stress are more common early life traumas affiliated with later mental illness and often co-occur in affected individuals.^{45,46,47} However, details regarding infants' *in utero* environment are not described in these studies, nor are details regarding the mothers' genetic makeup, rearing history, or mental health status—all of which are far likely more important contributors to the development of mental illness than the postnatal manipulations imposed by these researchers. Additionally, while macaque social structure may be as complex as human social structure, it is decidedly different from that of most modern human societies. For example, it is typical for infant macaques to stay in *constant* physical contact with their mothers for their first month of life,⁴⁸ making even the briefest separation stressful for infants as well as chronic separation more detrimental than can be expected in humans in most cultures. Therefore, any applicability of this nonhuman primate model is likely to vary dramatically across different human cultures with different social structures and traditional rearing practices. Even the “typical” mother-reared infants who are used as a control group in most of these experiments spend much of their time in barren, metal cages, and are subject to constant experimental testing, requiring multiple separations from their mother, and involving stress and/or fear-inducing tests.^{22,49} These living conditions and frequent maternal separations likely impact the natural infant-mother behavior that would occur in the wild, and as reviewed below, increase the stress levels and mental health of all animals included in the study. The mother-reared infants cannot provide an accurate example of “typical” or “healthy” development for any species, and the additional stress of laboratory conditions confound the experimental stressors introduced in maternally deprived animals. Therefore, these studies using a “well controlled” nonhuman primate model fail to properly model the complex relationship between genes, early life experience, and mental illness in the human population. The evidence of this fact is that, collectively, the project has not resulted in any new treatments for human mental illness. Almost four decades ago, Stephen Suomi himself acknowledged the limited applicability that his monkey experiments would have to human mental illness. In 1977, he wrote,

“...whether actual data obtained from nonhuman primates have added measurably to our understanding of human development is another matter....such cases are relatively rare. Most monkey data that readily generalize to humans have not uncovered new facts about human behavior; rather, they have only verified principles that have already been formulated from previous human data” (pg 203).⁵⁰

Existing clinical research and nonanimal methodologies readily available

The principal investigators on the aforementioned projects contend that controlled studies of gene-environment interactions in humans are ethically and practically untenable. However, this contention is

inaccurate. Numerous large-scale epidemiological studies in humans have documented the effects of early life stress,^{51,52,53} genetic risk,^{54,55,56} and gene-environment interactions,^{57,58,59,60,61,62} on abnormal social, emotional, and behavioral development. These studies include investigating the contribution of both genes and the environment in the development of mood disorders,^{57,58} addiction,⁶³ depression,⁶⁴ and altered brain structure and function^{65,66,67} in humans.

Recent human studies have also begun to unlock the complex biological and molecular mechanisms that underlie these gene and environmental interactions.^{65,68,69,70} For example, McGowan et al.⁶⁵ and Klengel et al.⁷⁰ studied the interaction between early childhood trauma and genetic variation on gene transcription in the brains of humans. Similarly, in a large-scale study of nearly 200 individuals, Buchman et al.⁶⁹ tested the interaction between early-life psychosocial adversity, genetic make-up, and plasma levels of brain-derived neurotrophic factor, critical for brain development and plasticity. DNA methylation, studied in the brain tissue of monkeys killed in the NIH studies, can be non-invasively measured in monocytes and T-cells and correlated with neurotransmitter synthesis using positron emission tomography *in vivo* in humans, a technique recently used to determine the relationship between childhood aggression, DNA methylation, and serotonergic function in humans.⁶⁸ Postmortem studies using brain tissue from humans at different stages of development⁷¹ as well as those from individuals suffering from or carrying genes associated with autism,^{72,73} depression,⁷⁴ and schizophrenia^{75,76} have identified critical differences in gene expression across age, species, and clinical populations. These groundbreaking studies have already begun detailing genetic *and* epigenetic effects on human brain structure, function, and development in humans suffering from mental illness—details not attainable from animal models.

Additionally, the mood-altering effects of the type of drugs being tested by the NIMH Non-Human Primate Core, including fluoxetine,⁷⁷ oxytocin,⁷⁸ diazepam,⁷⁹ and dopaminergic and serotonergic drugs such as raclopride and buspirone,^{80,81} are already well documented in humans suffering from mental illness. These studies have been conducted with healthy volunteers,^{82,83,84} children,^{75,85,86} and patients with mental illness.^{87,88} The impact that these drugs have on brain structure and function have also been evaluated in human volunteers,^{78,89,90} and their neural mechanisms in healthy and ill children and adults are already well delineated.^{91,92,93,94}

Impact on animal welfare

The physical and psychological harms of confining primates and other animals in laboratories and subjecting them to routine and experimental procedures are well established.^{95,96} Primates experience increased stress from common laboratory procedures such as cage cleaning,⁹⁷ physical examination,⁹⁸ blood draws,⁹⁹ and restraint.¹⁰⁰ The mere physical presence of human experimenters and technicians increases stress in primates.^{101,102} Numerous studies have demonstrated that even minor changes in primates' captive environment, including temporary changes in cage size or location, increase stress levels.^{103,104} It is not surprising that decreased immune system functioning¹⁰⁵ and increased self-injurious behavior are common in primates in laboratories.^{106,107}

Specific to the experiments in question, the intention of these projects is to create, psychological illness in primates. The numerous long-term negative outcomes of these motherless rearing conditions on monkeys have been well established for decades: mother-deprived infants exhibit excessive fearfulness and/or aggression,⁴⁸ produce excess stress hormones,¹⁰⁸ and frequently rank at the bottom of the social dominance hierarchy.⁴⁸ They exhibit motor stereotypies indicative of frustration and stress,¹⁰⁹ abnormal sleep patterns,¹¹⁰ increased susceptibility to alcohol abuse,¹¹¹ and increased startle and stress responses to threatening stimuli.¹¹² Maternal deprivation affects serotonin pathway function^{113,114} and

cerebral blood flow¹¹⁵ and alters levels of brain-derived neurotrophic factor and nerve growth factor critical for normal brain function¹¹⁶ and has long-term effects on brain morphology.¹¹⁷ Both spontaneous and selectively bred genetic variations in the macaques interact with adverse rearing conditions, often exacerbating the already profoundly negative effects of adverse rearing.^{118,119,120} Moreover, several of the procedures in this protocol, including the “Auditory Reflex,” “Response to Novelty,” and “Human Intruder” tests, require infants to frequently be restrained and subjected to stress and fear-inducing procedures for several hours a day, causing acute distress to the animals during the experimental trials. The repeated restraint and social isolation, repeated exposure to startling sounds and frightening situations, and repeated blood draws, spinal taps, drug injections, and brain imaging procedures cause more than temporary or minimal distress to the animals and take an enormous toll on the psychological well being of these animals.

Given the extensive, long-term psychological and physical harm caused by maternal deprivation, and the extensive distress the individual test procedures caused to animals included in this protocol, it would only be appropriate to classify these protocols as “USDA Column D” (“Pain or Distress Relieved By Appropriate Measures”) and/or “USDA Column E” (“unrelieved pain or distress”). However, the Animal Study Proposal (ASP 11-043) describing these procedures was classified as “USDA Column C,” indicating that the animals would suffer only “minimal or no pain or distress.” The investigators own research, cited above, document that this classification is inappropriate for this series of experiments. The ASP for these protocols was approved as “Column C” on May 02, 2011, and then again on April 16, 2014, through April 16, 2017. As a result of this misclassification, members of the Animal Care and Use Committee were not able to make an appropriate cost benefit analysis before approving the proposal. Moreover, investigators were not required to search for alternative, nonanimal methods for their studies—methods that, as described above, are readily available.

Additional independent review

To extend the depth of our analysis of these experiments, we have consulted with independent subject matter experts in the fields of mental health, medicine, anthropology, and primatology (they were not compensated in any way by PETA). Their concerns, which they have provided to PETA in writing, are as follows:

· Extensive experiments proving the damaging effects of maternal deprivation and isolation were carried out on rhesus monkeys by Harry Harlow and his students in the 1950s, 1960s and 1970s. And even after proof had been obtained, Harlow continued to devise ever more stressful situations... These experiments, getting more and more extreme, were unbelievably cruel. Nevertheless researchers continued working in this field after Harlow's death, and continue to do so today. ...It is my understanding that monkeys are being subjected to what I consider inhumane experiments at a laboratory in Maryland that is, to some extent, funded by public money. I was shown a video in which infant monkeys were taking part in experiments which I considered extremely cruel and unacceptable.....I am shocked and saddened that this is so.

Dr. Jane Goodall, DBE
Founder, the Jane Goodall Institute
UN Messenger of Peace

· *“Given the current status and progress of the research (as assessed via the published literature), I can no longer see a potential benefit from such experimentation as is ongoing currently. I cannot consider the depicted experiments, designed to create and study psychopathology in monkeys, to be a valuable undertaking that will likely contribute to the health and well being of humans.....From the methodologies described in the proposals and articles and the written and visual documentation provided by PETA of actual laboratory procedures and activities, it is my assessment that the monkeys used in these experiments experience substantial psychological (and likely physiological) harm and that there is no current evidence that there will be any results from the studies that move our understanding of human psychopathology forward.”*

Dr. Agustín Fuentes
Professor and Chair, Department of Anthropology
University of Notre Dame

· *In the past, I conducted experiments on the impact of social deprivation on monkey intelligence and abnormal social behavior. I eventually chose to leave that area of research because I came to believe that those models did not accurately represent the development and presentation of human mental illness. I came to the view that those models could not adequately inform innovative directions for successful clinical intervention to justify the costs in suffering and pain. I see nothing to alter that view with respect to the program of primate deprived early experience research currently being conducted at the NIH.*

Dr. John P. Gluck
Emeritus Professor of Psychology, University of New Mexico
Research Professor, Kennedy Institute of Ethics, Georgetown University

· *For the past three decades Suomi’s group has deprived hundreds of infant macaques of maternal contact to cause them to suffer from a range of severe and persistent cognitive, social, emotional, and physical deficits. There is no compelling evidence that these studies are now or have ever been beneficial to humans. This line of outdated and irrelevant research has unfortunately been allowed to proceed without proper scientific and ethical scrutiny by the NIH and the research community.*

Dr. Lori Marino
Executive Director, The Kimmela Center for Animal Advocacy
Former Senior Lecturer in Neuroscience and Animal Behavior, Emory University

· *“The cause of mental illness in humans is unknown, but it is clearly complex and multifactorial. Some genetic studies are promising. Abusing monkeys, however, won't get us any closer to that understanding.”*

Dr. Jaymie Shanker
Psychiatrist

· *“Taken as a group and without exception, these experiments are cruel, plunging infant monkeys into hellish conditions that they can neither control nor escape from. Ethically and morally, they have no place in science today. The cost to these animals is far too high. As we*

have seen, it is not as if the experiments lead to an earth-shattering breakthrough that could, in some moral calculus (though not PETA's and not mine), give us reason to think the cost was remotely worth it. This lack of justification is particularly true given the myriad of human-based research methodologies available to study the environmental, genetic, and social causes of mental illness as well as the fact that these experiments on monkeys often seek to replicate knowledge already ascertained in humans."

Dr. Barbara J. King
Chancellor Professor of Anthropology
College of William and Mary

· "The scientific objections to continuing this research are immediately obvious. If the goal is to model neuropathologic/neurophysiologic substrates of human psychiatric diseases, then these efforts are hopelessly crude and antiquated, having long been superseded by in vivo neuroimaging studies of human patients with the psychiatric diseases of interest. Simply conduct a search in PubMed on any psychiatric diagnosis, such as psychopathic personality disorder, depression, schizophrenia, and a host of others, and you will find dozens of current, sophisticated, state-of-the-art neuroimaging studies comparing brain structure and function in patients and controls, clearly delineating structural and functional abnormalities in human patients. These patients, along with their early life experiences, genetic make-up, and medical histories, can be followed longitudinally to evaluate illness etiology and treatment efficacy. Modern research methodology has also allowed investigators to measure the separate and interacting contribution of genes and early environmental stress in the development and neural substrates of mental illnesses in humans. Postmortem studies of human brain tissue from individuals with mental illnesses or individuals carrying risk-alleles associated with psychiatric diseases are far better methods for clarifying the molecular etiologies of these complex ailments.... If the goal of the infant monkey psychological trauma experiments is not to eventually improve our understanding of human psychiatric diseases—as the above cited imaging, genetic, and epidemiological studies are already doing—then in the zero sum game of research funding, the National Institutes of Health (presumably referring to human health) should have nothing to do with them."

Dr. Lawrence A. Hansen
Professor and Researcher, Department of Pathology
University of California, San Diego, School of Medicine

· "It is not surprising that monkeys reared under such adverse conditions at the NIH are physically, mentally, and emotionally unwell. However, despite the outcome being known, it is surprising that experiments in which these animals are deliberately subjected to extreme stress are allowed to continue. Moreover, monkeys are not humans, so any experimental findings that are true of monkeys would not necessarily be true of humans. If the researchers who are performing these experiments wish to argue that the monkeys are similar enough to humans in terms of emotional development that studies done on them can be applied to human development, then they must acknowledge that they are performing studies that cause intense pain and terror to their subjects, much as any human would experience intense pain and terror were these experiments performed on humans..... The American Psychological Association should not permit these experiments, which I believe are in violation of several sections of the APA Guidelines for Ethical Conduct in the Care and Use of Nonhuman Animals in Research.

Specifically, Guideline I (2) states that “[T]he scientific purpose of the research should be of sufficient potential significance to justify the use of nonhuman animals” and notes that “psychologists should act on the assumption that procedures that are likely to produce pain in humans may also do so in other animals.” Yet, in a 2014 paper published in The American Journal of Psychiatry, the experimenters acknowledge that their anti-depressant experiments on monkeys cannot be applied to humans, that maternal deprivation studies on monkeys have never been confirmed as an effective way to test the efficacy of drug treatments for human mental illness, and that the only way to test treatments for human psychological disorders is in humans.”

Dr. Michael Radkowsky
Clinical Psychologist

· “If these experiments are meant to parallel or predict the psychopathy and mental illness of human infants in the care of negligent, absent, and/or abusive mothers, they fail profoundly. Contrived maternal deprivation, chronic exposure to stressful experimental paradigms, confinement, and social isolation in laboratory settings do not parallel the types of early stressors experienced by most human mental illness sufferers. These laboratory versions of early-life adversity are too routinized and methodical to be representative of any real-world experiences faced by humans. The circumstances surrounding physical, social, emotional, and cognitive development in human beings is multifaceted and more complicated than those that can be imposed on infant monkeys reared in a laboratory. Good, creative research either cleverly sets up situations that allow behavioral and biological responses of interest to occur naturally, or it takes the form of field studies to observe real-world dynamics in a natural setting. The NIH experiments depicted on video include constraining infants in small cages and startling them with loud noises, trapping infants and then threatening them with human experimenters, or caging them with a drugged, unresponsive mother. These procedures do not accurately or creatively replicate the stressful situations believed to precipitate mental illness in humans.”

Dr. Nora J. Johnson
Clinical Psychologist
University of Pennsylvania Health System

· “I do not consider the depicted experiments, designed to create and study psychopathology in monkeys, to be a valuable undertaking that will likely contribute to the health and well-being of humans. Rather, the causes and manifestations of mental illness in humans are most effectively researched without the use of animals.”

Dr. Tara West
Professor, Social Psychologist
CUNY School of Professional Studies

Conclusion

In a recent paper discussing the inadequacy of regulations governing experimentation on animals, bioethicist Dr. David Wendler of the NIH’s Clinical Center called for greater restrictions on the use of primates in experiments, noting that existing regulations “do not mandate that the risks to which nonhuman primates are exposed must be justified by the value of the study in question.”¹²¹

- 1 Ferrari, P. F., Vanderwert, R. E., Paukner, A., Bower, S., Suomi, S. J., & Fox, N. A. (2012). Distinct EEG amplitude suppression to facial gestures as evidence for a mirror mechanism in newborn monkeys. **Journal of Cognitive Neuroscience**, *24*(5), 1165-1172.
- 2 Vanderwert, R. E., Ferrari, P. F., Paukner, A., Bower, S. B., Fox, N. A., & Suomi, S. J. (2012). Spectral characteristics of the newborn rhesus macaque EEG reflect functional cortical activity. **Physiology & Behavior**, *107*(5), 787-791.
- Ferrari, P. F., Vanderwert, R. E., Paukner, A., Bower, S., Suomi, S. J., & Fox, N. A. (2012). Distinct EEG amplitude suppression to facial gestures as evidence for a mirror mechanism in newborn monkeys. **Journal of Cognitive Neuroscience**, *24*(5), 1165-1172.
- 5 Vanderwert, R. E., Ferrari, P. F., Paukner, A., Bower, S. B., Fox, N. A., & Suomi, S. J. (2012). Spectral characteristics of the newborn rhesus macaque EEG reflect functional cortical activity. **Physiology & Behavior**, *107*(5), 787-791.
- e, L., Alleva, E., & Suomi, S. J. (2009). Early life stress as a risk factor for mental health: role of neurotrophins from rodents to non-human primates. **Neuroscience & Biobehavioral Reviews**, *33*(4), 573-585.
- 7 Dettmer, A. M., Novak, M. F., Novak, M. A., Meyer, J. S., & Suomi, S. J. (2009). Hair cortisol predicts object permanence performance in infant rhesus macaques (*Macaca mulatta*). **Developmental Psychobiology**, *51*(8), 706-713.
- 8 Newman, T. K., Syagailo, Y. V., Barr, C. S., Wendland, J. R., Champoux, M., Graessle, M., ... & Lesch, K. P. (2005). Monoamine oxidase A gene promoter variation and rearing experience influences aggressive behavior in rhesus monkeys. **Biological Psychiatry**, *57*(2), 167-172.
- 9 Spinelli, S., Schwandt, M. L., Lindell, S. G., Newman, T. K., Heilig, M., Suomi, S. J., ... & Barr, C. S. (2007). Association between the recombinant human serotonin transporter linked promoter region polymorphism and behavior in rhesus macaques during a separation paradigm. **Development and Psychopathology**, *19*(04), 977-987.
- 10 Erickson, K., Gabry, K. E., Schulkin, J., Gold, P., Lindell, S., Higley, J. D., ... & Suomi, S. J. (2005). Social withdrawal behaviors in nonhuman primates and changes in neuroendocrine and monoamine concentrations during a separation paradigm. **Developmental Psychobiology**, *46*(4), 331-339.
- 11 Barr, C. S., Newman, T. K., Shannon, C., Parker, C., Dvoskin, R. L., Becker, M. L., ... & Higley, J. D. (2004). Rearing condition and rh5-HTTLPR interact to influence limbic-hypothalamic-pituitary-adrenal axis response to stress in infant macaques. **Biological Psychiatry**, *55*(7), 733-738.
- 12 Champoux, M., Bennett, A., Shannon, C., Higley, J. D., Lesch, K. P., & Suomi, S. J. (2002). Serotonin transporter gene polymorphism, differential early rearing, and behavior in rhesus monkey neonates. **Molecular Psychiatry**, *7*(10), 1058-1063.
- 13 Higley, J. D., Suomi, S. J., & Linnoila, M. (1991). CSF monoamine metabolite concentrations vary according to age, rearing, and sex, and are influenced by the stressor of social separation in rhesus monkeys. **Psychopharmacology**, *103*(4), 551-556.
- 14 Westergaard, G. C., Mehlman, P. T., Suomi, S. J., & Higley, J. D. (1999). CSF 5-HIAA and aggression in female macaque monkeys: species and interindividual differences. **Psychopharmacology**, *146*(4), 440-446.
- 15 Higley, J. D., Thompson, W. W., Champoux, M., Goldman, D., Hasert, M. F., Kraemer, G. W., ... & Linnoila, M. (1993). Paternal and maternal genetic and environmental contributions to cerebrospinal fluid monoamine metabolites in rhesus monkeys (*Macaca mulatta*). **Archives of General Psychiatry**, *50*(8), 615-623.
- 16 Higley, J. D., Suomi, S. J., & Linnoila, M. (1992). A longitudinal assessment of CSF monoamine metabolite and plasma cortisol concentrations in young rhesus monkeys. **Biological Psychiatry**, *32*(2), 127-145.
- For decades the NIH has continued to review, approve, fund, and conduct the aforementioned studies that deliberately and repeatedly inflict severe and chronic harm to monkeys, are often not at all designed to help humans, or have extremely limited potential to elucidate the complex etiology of human mental illness and have not improved our treatments of these illnesses or human health in general.
- These experiments represent an enormous financial burden to taxpayers, particularly as there are a myriad of accessible, humane research methodologies that are more directly applicable to mental illness and its treatment. Continuing to fund this suite of projects appears to be both scientifically and ethically unjustifiable.

References

- 17 Paukner, A., Huntsberry, M. E., & Suomi, S. J. (2010). Visual discrimination of male and female faces by infant rhesus macaques. **Developmental Psychobiology**, **52**(1), 54-61.
- 18 Paukner, A., Ferrari, P. F., & Suomi, S. J. (2011). Delayed imitation of lipsmacking gestures by infant rhesus macaques (*Macaca mulatta*). **PLoS One**, **6**(12), e28848.
- 19 Little, A. C., Paukner, A., Woodward, R. A., & Suomi, S. J. (2012). Facial asymmetry is negatively related to condition in female macaque monkeys. **Behavioral Ecology and Sociobiology**, **66**(9), 1311-1318.
- 20 Bower, S., Suomi, S. J., & Paukner, A. (2012). Evidence for kinship information contained in the rhesus macaque (*Macaca mulatta*) face. **Journal of Comparative Psychology**, **126**(3), 318.
- 21 Paukner, A., Bower, S., Simpson, E. A., & Suomi, S. J. (2013). Sensitivity to first-order relations of facial elements in infant rhesus macaques. **Infant and Child Development**, **22**(3), 320-330.
- 22 Zhang, B., Suarez-Jimenez, B., Hathaway, A., Waters, C., Vaughan, K., Noble, P. L., ... & Nelson, E. E. (2012). Developmental changes of rhesus monkeys in response to separation from the mother. **Developmental Psychobiology**, **54**(8), 798-807.
- 23 Suarez-Jimenez B., Hathaway, A., Waters, C., Vaughan, K., Suomi, S. J., Noble, P. L., ... & Nelson, E. E. (2013). Effect of mother's dominance rank on offspring temperament in infant rhesus monkeys (*Macaca mulatta*). **American Journal of Primatology**, **75**(1), 65-73.
- 24 Shrestha, S. S., Nelson, E. E., Liow, J. S., Gladding, R., Lyoo, C. H., Noble, P. L., ... & Innis, R. B. (2014). Fluoxetine administered to juvenile monkeys: effects on the serotonin transporter and behavior. **American Journal of Psychiatry**, **171**(3), 323-331.
- 25 Spinelli, S., Schwandt, M. L., Lindell, S. G., Heilig, M., Suomi, S. J., Higley, J. D., ... & Barr, C. S. (2012). The serotonin transporter gene linked polymorphic region is associated with the behavioral response to repeated stress exposure in infant rhesus macaques. **Development and Psychopathology**, **24**(01), 157-165.
- 26 Lindell, S. G., Schwandt, M. L., Sun, H., Sparenborg, J. D., Björk, K., Kasckow, J. W., ... & Barr, C. S. (2010). Functional NPY variation as a factor in stress resilience and alcohol consumption in rhesus macaques. **Archives of General Psychiatry**, **67**(4), 423-431.
- 27 Schwandt, M. L., Lindell, S. G., Higley, J. D., Suomi, S. J., Heilig, M., & Barr, C. S. (2011). OPRM1 gene variation influences hypothalamic-pituitary-adrenal axis function in response to a variety of stressors in rhesus macaques. **Psychoneuroendocrinology**, **36**(9), 1303-1311.
- 28 Barr, C. S., Becker, M. L., Suomi, S. J., & Higley, J. D. (2003). Relationships among CSF monoamine metabolite levels, alcohol sensitivity, and alcohol-related aggression in rhesus macaques. **Aggressive Behavior**, **29**(4), 288-301.
- 29 Barr, C. S., Newman, T. K., Lindell, S., Shannon, C., Champoux, M., Lesch, K. P., ... & Higley, J. D. (2004). Interaction between serotonin transporter gene variation and rearing condition in alcohol preference and consumption in female primates. **Archives of General Psychiatry**, **61**(11), 1146-1152.
- 30 Barr, C. S., Dvoskin, R. L., Gupte, M., Sommer, W., Sun, H., Schwandt, M. L., ... & Heilig, M. (2009). Functional CRH variation increases stress-induced alcohol consumption in primates. **Proceedings of the National Academy of Sciences**, **106**(34), 14593-14598.
- 31 Schwandt, M. L., Lindell, S. G., Chen, S., Higley, J. D., Suomi, S. J., Heilig, M., & Barr, C. S. (2010). Alcohol response and consumption in adolescent rhesus macaques: life history and genetic influences. **Alcohol**, **44**(1), 67-80.
- 32 Alexander, G. M., Graef, J. D., Hammarback, J. A., Nordskog, B. K., Burnett, E. J., Daunais, J. B., ... & Godwin, D. W. (2012). Disruptions in serotonergic regulation of cortical glutamate release in primate insular cortex in response to chronic ethanol and nursery rearing. **Neuroscience**, **207**, 167-181.
- 33 Huggins, K. N., Mathews, T. A., Locke, J. L., Szeliga, K. T., Friedman, D. P., Bennett, A. J., & Jones, S. R. (2012). Effects of early life stress on drinking and serotonin system activity in rhesus macaques: 5-hydroxyindoleacetic acid in cerebrospinal fluid predicts brain tissue levels. **Alcohol**, **46**(4), 371-376.
- 34 Provençal, N., Suderman, M. J., Guillemin, C., Massart, R., Ruggiero, A., Wang, D., ... & Szyf, M. (2012). The signature of maternal rearing in the methylome in rhesus macaque prefrontal cortex and T cells. **Journal of Neuroscience**, **32**(44), 15626-15642.
- 35 Enard, W., Khaitovich, P., Klose, J., Zöllner, S., Heissig, F., Giavalisco, P., ... & Pääbo, S. (2002). Intra- and interspecific variation in primate gene expression patterns. **Science**, **296**(5566), 340-343.
- 36 Cáceres, M., Lachuer, J., Zapala, M. A., Redmond, J. C., Kudo, L., Geschwind, D. H., ... & Barlow, C. (2003). Elevated gene expression levels distinguish human from non-human primate brains. **Proceedings of the National Academy of Sciences**, **100**(22), 13030-13035.
- 37 Shi, L., Li, M., Lin, Q., Qi, X., & Su, B. (2013). Functional divergence of the brain-size regulating gene MCPH1 during primate evolution and the origin of humans. **BMC Biology**, **11**(1), 62.

³⁸ Muntané, G., Horvath, J. E., Hof, P. R., Ely, J. J., Hopkins, W. D., Raghanti, M. A., ... & Sherwood, C. C. (2014). Analysis of synaptic gene expression in the neocortex of primates reveals evolutionary changes in glutamatergic neurotransmission. **Cerebral Cortex**, **bht354**.

- 39 Balsters, J. H., Cussans, E., Diedrichsen, J., Phillips, K. A., Preuss, T. M., Rilling, J. K., & Ramnani, N. (2010). Evolution of the cerebellar cortex: the selective expansion of prefrontal-projecting cerebellar lobules. **Neuroimage**, **49(3)**, 2045-2052.
- 40 Fu, X., Giavalisco, P., Liu, X., Catchpole, G., Fu, N., Ning, Z. B., ... & Khaitovich, P. (2011). Rapid metabolic evolution in human prefrontal cortex. **Proceedings of the National Academy of Sciences**, **108(15)**, 6181-6186.
- 41 Hecht, E. E., Gutman, D. A., Preuss, T. M., Sanchez, M. M., Parr, L. A., & Rilling, J. K. (2013). Process versus product in social learning: comparative diffusion tensor imaging of neural systems for action execution–observation matching in macaques, chimpanzees, and humans. **Cerebral Cortex**, **23(5)**, 1014-1024.
- 42 Rilling, J. K. (2014). Comparative primate neuroimaging: insights into human brain evolution. **Trends in Cognitive Sciences**, **18(1)**, 46-55.
- 43 Geschwind, D. H., & Rakic, P. (2013). Cortical evolution: judge the brain by its cover. **Neuron**, **80(3)**, 633-647.
- 44 Sakai, T., Matsui, M., Mikami, A., Malkova, L., Hamada, Y., Tomonaga, M., ... & Matsuzawa, T. (2013). Developmental patterns of chimpanzee cerebral tissues provide important clues for understanding the remarkable enlargement of the human brain. **Proceedings of the Royal Society B: Biological Sciences**, **280(1753)**.
- 45 Green, J. G., McLaughlin, K. A., Berglund, P. A., Gruber, M. J., Sampson, N. A., Zaslavsky, A. M., et al. (2010). Childhood adversities and adult psychiatric disorders in the national comorbidity survey replication I: associations with first onset of DSM-IV disorders. **Archives of General Psychiatry**, **67(2)**, 113-123.
- 46 McLaughlin, K. A., Green, J. G., Gruber, M. J., Sampson, N. A., Zaslavsky, A. M., & Kessler, R. C. (2010). Childhood adversities and adult psychiatric disorders in the national comorbidity survey replication II: associations with persistence of DSM-IV disorders. **Arch Gen Psychiatry**, **67(2)**, 124-132.
- 47 McLaughlin, K. A., Gadermann, A. M., Hwang, I., Sampson, N. A., Al-Hamzawi, A., Andrade, L. H., ... & Kessler, R. C. (2012). Parent psychopathology and offspring mental disorders: results from the WHO World Mental Health Surveys. **British Journal of Psychiatry**, **200(4)**, 290-299.
- 48 Suomi, S. J. (1997). Early determinants of behavior: evidence from primate studies. **Br Med Bull**, **53**, 170-184.
- 49 Spinelli, S., Schwandt, M. L., Lindell, S. G., Heilig, M., Suomi, S. J., Higley, J. D., ... & Barr, C. S. (2012). The serotonin transporter gene linked polymorphic region is associated with the behavioral response to repeated stress exposure in infant rhesus macaques. **Development and Psychopathology**, **24(01)**, 157-165.
- 50 Suomi, S. J. (1976). Mechanisms underlying social development: a re-examination of mother-infant interactions in monkeys. In: *Minnesota Symposium on child psychology* (Vol. 10, pp. 201-228).
- 51 Neigh, G. N., Gillespie, C. F., & Nemeroff, C. B. (2009). The neurobiological toll of child abuse and neglect. **Trauma, Violence, & Abuse**, **10(4)**, 389-410.
- 52 Greeson, J. K., Briggs, E. C., Kisiel, C. L., Layne, C. M., Ake III, G. S., Ko, S. J., ... & Fairbank, J. A. (2011). Complex trauma and mental health in children and adolescents placed in foster care: findings from the National Child Traumatic Stress Network. **Child Welfare**, **90(6)**.
- 53 Tottenham, N. (2012). Risk and developmental heterogeneity in previously institutionalized children. **Journal of Adolescent Health**, **51(2)**, S29-S33.
- 54 Grabe, H. J., Schwahn, C., Appel, K., Mahler, J., Schulz, A., Spitzer, C., ... & Völzke, H. (2010). Childhood maltreatment, the corticotropin-releasing hormone receptor gene and adult depression in the general population. **American Journal of Medical Genetics Part B: Neuropsychiatric Genetics**, **153(8)**, 1483-1493.
- 55 Grabe, H. J., Schwahn, C., Mahler, J., Appel, K., Schulz, A., Spitzer, C., ... & Völzke, H. (2012). Genetic epistasis between the brain-derived neurotrophic factor Val66Met polymorphism and the 5-HTT promoter polymorphism moderates the susceptibility to depressive disorders after childhood abuse. **Progress in Neuro-Psychopharmacology and Biological Psychiatry**, **36(2)**, 264-270.
- 56 Caldwell, W., McInnis, O. A., McQuaid, R. J., Liu, G., Stead, J. D., Anisman, H., & Hayley, S. (2013). The role of the Val66Met polymorphism of the brain derived neurotrophic factor gene in coping strategies relevant to depressive symptoms. **PloS One**, **8(6)**, e65547.
- 57 Taylor, S. E., Way, B. M., Welch, W. T., Hilmert, C. J., Lehman, B. J., & Eisenberger, N. I. (2006). Early family environment, current adversity, the serotonin transporter promoter polymorphism, and depressive symptomatology. **Biological Psychiatry**, **60(7)**, 671-676.
- 58 Stein, M. B., Schork, N. J., & Gelernter, J. (2008). Gene-by-environment (serotonin transporter and childhood maltreatment) interaction for anxiety sensitivity, an intermediate phenotype for anxiety disorders. **Neuropsychopharmacology**, **33(2)**, 312-319.
- 59 Armbruster, D., Mueller, A., Strobel, A., Lesch, K. P., Brocke, B., & Kirschbaum, C. (2012). Children under stress–COMT genotype and stressful life events predict cortisol increase in an acute social stress paradigm. **International Journal of Neuropsychopharmacology**, **15(09)**, 1229-1239.
- 60 Carver, C. S., Johnson, S. L., Joormann, J., Kim, Y., & Nam, J. Y. (2011). Serotonin transporter polymorphism interacts with childhood adversity to predict aspects of impulsivity. **Psychological Science**, **22(5)**, 589-595.

- 61 Drury, S. S., Gleason, M. M., Theall, K. P., Smyke, A. T., Nelson, C. A., Fox, N. A., & Zeanah, C. H. (2012). Genetic sensitivity to the caregiving context: The influence of 5httlpr and BDNF val66met on indiscriminate social behavior. **Physiology & Behavior**, **106(5)**, 728-735.
- 62 Uher, R., Caspi, A., Houts, R., Sugden, K., Williams, B., Poulton, R., & Moffitt, T. E. (2011). Serotonin transporter gene moderates childhood maltreatment's effects on persistent but not single-episode depression: replications and implications for resolving inconsistent results. **Journal of Affective Disorders**, **135(1)**, 56-65.
- 63 Enoch, M. A. (2011). The role of early life stress as a predictor for alcohol and drug dependence. **Psychopharmacology**, **214(1)**, 17-31.
- 64 Aguilera, M., Arias Sampérez, B., Wichers, M., Barrantes Vidal, N., Moya Higuera, J., Villa Martín, E., ... & Fañanás Saura, L. (2009). Early adversity and 5-HTT-BDNF genes: new evidences of gene-environment interactions on depressive symptoms in a general population. **Psychological Medicine** ISSN, **39(9)**, 1425-1432.
- 65 McGowan, P. O., Sasaki, A., D'Alessio, A. C., Dymov, S., Labonté, B., Szyf, M., ... & Meaney, M. J. (2009). Epigenetic regulation of the glucocorticoid receptor in human brain associates with childhood abuse. **Nature Neuroscience**, **12(3)**, 342-348.
- 66 Sheridan, M. A., Fox, N. A., Zeanah, C. H., McLaughlin, K. A., & Nelson, C. A. (2012). Variation in neural development as a result of exposure to institutionalization early in childhood. **Proceedings of the National Academy of Sciences**, **109(32)**, 12927-12932.
- 67 Frodl, T., & O'Keane, V. (2013). How does the brain deal with cumulative stress? A review with focus on developmental stress, HPA axis function and hippocampal structure in humans. **Neurobiology of Disease**, **52**, 24-37.
- 68 Wang D., Szyf M., Benkelfat C., Provencal N., Turecki G., et al. (2012). Peripheral SLC6A4 DNA methylation is associated with in vivo measures of human brain serotonin synthesis and childhood physical aggression. **PLoS One** **7(6)**.
- 69 Buchmann, A. F., Hellweg, R., Rietschel, M., Treutlein, J., Witt, S. H., Zimmermann, U. S., ... & Deuschle, M. (2013). BDNF Val 66 Met and 5-HTTLPR genotype moderate the impact of early psychosocial adversity on plasma brain-derived neurotrophic factor and depressive symptoms: A prospective study. **European Neuropsychopharmacology**, **23(8)**, 902-909.
- 70 Klengel, T., Mehta, D., Anacker, C., Rex-Haffner, M., Pruessner, J. C., Pariante, C. M., ... & Binder, E. B. (2013). Allele-specific FKBP5 DNA demethylation mediates gene-childhood trauma interactions. **Nature Neuroscience**, **16(1)**, 33-41.
- 71 Miller, J. A., Ding, S. L., Sunkin, S. M., Smith, K. A., Ng, L., Szafer, A., ... & Pletikos, M. (2014). Transcriptional landscape of the prenatal human brain. **Nature**, **508**, 199-206.
- 72 James, S. J., Shpyleva, S., Melnyk, S., Pavliv, O., & Pogribny, I. P. (2013). Complex epigenetic regulation of Engrailed-2 (EN-2) homeobox gene in the autism cerebellum. **Translational psychiatry**, **3(2)**, e232.
- 73 Stoner, R., Chow, M. L., Boyle, M. P., Sunkin, S. M., Mouton, P. R., Roy, S., ... & Courchesne, E. (2014). Patches of disorganization in the neocortex of children with autism. **New England Journal of Medicine**, **370(13)**, 1209-1219.
- 74 Duric, V., Banasr, M., Stockmeier, C. A., Simen, A. A., Newton, S. S., Overholser, J. C., ... & Duman, R. S. (2013). Altered expression of synapse and glutamate related genes in post-mortem hippocampus of depressed subjects. **International Journal of Neuropsychopharmacology**, **16(01)**, 69-82.
- 75 Kunii, Y., Hyde, T. M., Ye, T., Li, C., Kolachana, B., Dickinson, D., ... & Lipska, B. K. (2014). Revisiting DARPP-32 in postmortem human brain: changes in schizophrenia and bipolar disorder and genetic associations with t-DARPP-32 expression. **Molecular Psychiatry**, **19(2)**, 192-199.
- 76 Wockner, L. F., Noble, E. P., Lawford, B. R., Young, R. M., Morris, C. P., Whitehall, V. L. J., & Voisey, J. (2014). Genome-wide DNA methylation analysis of human brain tissue from schizophrenia patients. **Translational Psychiatry**, **4(1)**, e339.
- 77 Emslie, G., Kennard, B., Mayes, T., Nightingale-Teresi, J., Carmody, T., Hughes, C., ... & Rintelmann, J. (2008). Fluoxetine versus placebo in preventing relapse of major depression in children and adolescents. **American Journal of Psychiatry**, **165(4)**, 459-467.
- 78 Mah, B. L., Van IJzendoorn, M. H., Smith, R., & Bakermans-Kranenburg, M. J. (2013). Oxytocin in postnatally depressed mothers: its influence on mood and expressed emotion. **Progress in Neuro-Psychopharmacology and Biological Psychiatry**, **40**, 267-272.
- 79 Delgado, V. B., Izquierdo, I., & Chaves, M. L. (2005). Differential effects of acute diazepam on emotional and neutral memory tasks in acutely hospitalized depressed patients. **Neuropsychiatr Dis Treat**, **1**, 269-275.

80 Anderer, P., Saletu, B., & Pascual-Marqui, R. D. (2000). Effect of the 5-HT_{1A} partial agonist buspirone on regional brain electrical activity in man: a functional neuroimaging study using low-resolution electromagnetic tomography (LORETA). **Psychiatry Research: Neuroimaging**, **100(2)**, 81-96.

81 Nordström, A. L., Farde, L., Wiesel, F. A., Forslund, K., Pauli, S., Halldin, C., & Uppfeldt, G. (1993). Central D₂-dopamine receptor occupancy in relation to antipsychotic drug effects: a double-blind PET study of schizophrenic patients. **Biological Psychiatry**, **33(4)**, 227-235.

- 82 Reynolds, B., Richards, J. B., Dassinger, M., & de Wit, H. (2004). Therapeutic doses of diazepam do not alter impulsive behavior in humans. **Pharmacology Biochemistry and Behavior**, **79(1)**, 17-24.
- 83 Di Simplicio, M., Massey-Chase, R., Cowen, P. J., & Harmer, C. J. (2009). Oxytocin enhances processing of positive versus negative emotional information in healthy male volunteers. **Journal of Psychopharmacology**, **23(3)**, 241-248.
- 84 McCabe, C., & Mishor, Z. (2011). Antidepressant medications reduce subcortical–cortical resting-state functional connectivity in healthy volunteers. **Neuroimage**, **57(4)**, 1317-1323.
- 85 Davari-Ashtiani, R., Shahrbabaki, M. E., Razjouyan, K., Amini, H., & Mazhabdar, H. (2010). Buspirone versus methylphenidate in the treatment of attention deficit hyperactivity disorder: a double-blind and randomized trial. **Child Psychiatry & Human Development**, **41(6)**, 641-648.
- 86 Gordon, I., Vander Wyk, B. C., Bennett, R. H., Cordeaux, C., Lucas, M. V., Eilbott, J. A., ... & Pelphey, K. A. (2013). Oxytocin enhances brain function in children with autism. **Proceedings of the National Academy of Sciences**, **110(52)**, 20953-20958.
- 87 Rickels, K., Downing, R., Schweizer, E., & Hassman, H. (1993). Antidepressants for the treatment of generalized anxiety disorder: a placebo-controlled comparison of imipramine, trazodone, and diazepam. **Archives of General Psychiatry**, **50(11)**, 884-895.
- 88 Guastella, A. J., Howard, A. L., Dadds, M. R., Mitchell, P., & Carson, D. S. (2009). A randomized controlled trial of intranasal oxytocin as an adjunct to exposure therapy for social anxiety disorder. **Psychoneuroendocrinology**, **34(6)**, 917-923.
- 89 Del-Ben, C. M., Ferreira, C. A., Sanchez, T. A., Alves-Neto, W. C., Guapo, V. G., de Araujo, D. B., & Graeff, F. G. (2012). Effects of diazepam on BOLD activation during the processing of aversive faces. **Journal of Psychopharmacology**, **26(4)**, 443-451.
- 90 Domes, G., Heinrichs, M., Gläscher, J., Büchel, C., Braus, D. F., & Herpertz, S. C. (2007). Oxytocin attenuates amygdala responses to emotional faces regardless of valence. **Biological Psychiatry**, **62(10)**, 1187-1190.
- 91 Ceccarini, J., Vrieze, E., Koole, M., Muylle, T., Bormans, G., Claes, S., & Van Laere, K. (2012). Optimized in vivo detection of dopamine release using 18F-fallypride PET. **Journal of Nuclear Medicine**, **53(10)**, 1565-1572.
- 92 Loane, C., & Politis, M. (2012). Buspirone: What is it all about? **Brain Research**, **1461**, 111-118.
- 93 Tamaji, A., Iwamoto, K., Kawamura, Y., Takahashi, M., Ebe, K., Kawano, N., ... & Ozaki, N. (2012). Differential effects of diazepam, tandospirone, and paroxetine on plasma brain-derived neurotrophic factor level under mental stress. **Human Psychopharmacology: Clinical and Experimental**, **27(3)**, 329-333.
- 94 Kanat, M., Heinrichs, M., & Domes, G. (2013). Oxytocin and the social brain: Neural mechanisms and perspectives in human research. **Brain Research**. [Epub ahead of print]
- 95 Balcombe, J. P., Barnard, N. D., & Sandusky, C. (2004). Laboratory routines cause animal stress. **Journal of the American Association for Laboratory Animal Science**, **43(6)**, 42-51.
- 96 Ferdowsian, H., & Merskin, D. (2012). Parallels in sources of trauma, pain, distress, and suffering in humans and nonhuman animals. **Journal of Trauma & Dissociation**, **13(4)**, 448-468.
- 97 Line, S.W., Morgan, K.N., Markowitz, H., Strong, S. (1989). Heart rate and activity of rhesus monkeys in response to routine events. **Laboratory Primate Newsletter**, **28(2)**, 1-4.
- 98 Golub, M. S., & Anderson, J. H. (1986). Adaptation of pregnant rhesus monkeys to short-term chair restraint. **Laboratory Animal Science**, **36(5)**, 507-511.
- 99 Gordon, T. P., Gust, D. A., Wilson, M. E., Ahmed-Ansari, A., Brodie, A. R., & McClure, H. M. (1992). Social separation and reunion affects immune system in juvenile rhesus monkeys. **Physiology & Behavior**, **51(3)**, 467-472.
- 100 Fuller, G. B., Hobson, W. C., Reyes, F. I., Winter, J. S. D., & Faiman, C. (1984). Influence of restraint and ketamine anesthesia on adrenal steroids, progesterone, and gonadotropins in rhesus monkeys. **Experimental Biology and Medicine**, **175(4)**, 487-490.
- 101 Barros, M., & Tomaz, C. (2002). Non-human primate models for investigating fear and anxiety. **Neuroscience & Biobehavioral Reviews**, **26(2)**, 187-201.
- 102 Suzuki, J., Ohkura, S., & Terao, K. (2002). Baseline and stress levels of cortisol in conscious and unrestrained Japanese macaques (*Macaca fuscata*). **Journal of Medical Primatology**, **31(6)**, 340-344.
- 103 Crockett, C. M., Shimoji, M., & Bowden, D. M. (2000). Behavior, appetite, and urinary cortisol responses by adult female pigtailed macaques to cage size, cage level, room change, and ketamine sedation. **American Journal of Primatology**, **52(2)**, 63-80.
- 104 Reinhardt, V., & Reinhardt, A. (2000). The lower row monkey cage: An overlooked variable in biomedical research. **Journal of Applied Animal Welfare Science**, **3(2)**, 141-149.

¹⁰⁵ Schapiro, S. J., Nehete, P. N., Perlman, J. E., & Sastry, K. J. (2000). A comparison of cell-mediated immune responses in rhesus macaques housed singly, in pairs, or in groups. **Applied Animal Behaviour Science**, **68(1)**, 67-84.

¹⁰⁶ Novak, M. A. (2003). Self-injurious behavior in rhesus monkeys: new insights into its etiology, physiology, and treatment. **American Journal of Primatology**, **59(1)**, 3-19.

- 107 Rommeck, I., Anderson, K., Heagerty, A., Cameron, A., & McCowan, B. (2009). Risk factors and remediation of self-injurious and self-abuse behavior in rhesus macaques. **Journal of Applied Animal Welfare Science**, **12(1)**, 61-72.
- 108 Dettmer, A. M., Novak, M. A., Suomi, S. J., & Meyer, J. S. (2012). Physiological and behavioral adaptation to relocation stress in differentially reared rhesus monkeys: hair cortisol as a biomarker for anxiety-related responses. **Psychoneuroendocrinology**, **37(2)**, 191-199.
- 109 Barr, C. S., Becker, M. L., Suomi, S. J., & Higley, J. D. (2003). Relationships among CSF monoamine metabolite levels, alcohol sensitivity, and alcohol-related aggression in rhesus macaques. **Aggressive Behavior**, **29(4)**, 288-301.
- 110 Barrett, C. E., Noble, P., Hanson, E., Pine, D. S., Winslow, J. T., & Nelson, E. E. (2009). Early adverse rearing experiences alter sleep-wake patterns and plasma cortisol levels in juvenile rhesus monkeys. **Psychoneuroendocrinology**, **34(7)**, 1029-1040.
- 111 Fahlke, C., Lorenz, J. G., Long, J., Champoux, M., Suomi, S. J., & Higley, J. D. (2000). Rearing experiences and stress-induced plasma cortisol as early risk factors for excessive alcohol consumption in nonhuman primates. **Alcoholism: Clinical and Experimental Research**, **24(5)**, 644-650.
- 112 Nelson, E. E., Herman, K. N., Barrett, C. E., Noble, P. L., Wojteczko, K., Chisholm, K., ... & Pine, D. S. (2009). Adverse rearing experiences enhance responding to both aversive and rewarding stimuli in juvenile rhesus monkeys. **Biological Psychiatry**, **66(7)**, 702-704.
- 113 Bennett, A. J., Lesch, K. P., Heils, A., Long, J. C., Lorenz, J. G., Shoaf, S. E., ... & Higley, J. D. (2002). Early experience and serotonin transporter gene variation interact to influence primate CNS function. **Molecular Psychiatry**, **7(1)**, 118-122.
- 114 Spinelli, S., Chefer, S., Carson, R. E., Jagoda, E., Lang, L., Heilig, M., ... & Stein, E. A. (2010). Effects of early-life stress on serotonin 1A receptors in juvenile rhesus monkeys measured by positron emission tomography. **Biological Psychiatry**, **67(12)**, 1146-1153.
- 115 Ichise, M., Vines, D. C., Gura, T., Anderson, G. M., Suomi, S. J., Higley, J. D., & Innis, R. B. (2006). Effects of early life stress on [11C] DASB positron emission tomography imaging of serotonin transporters in adolescent peer- and mother-reared rhesus monkeys. **Journal of Neuroscience**, **26(17)**, 4638-4643.
- 116 Cirulli, F., Francia, N., Branchi, I., Antonucci, M. T., Aloe, L., Suomi, S. J., & Alleva, E. (2009). Changes in plasma levels of BDNF and NGF reveal a gender-selective vulnerability to early adversity in rhesus macaques. **Psychoneuroendocrinology**, **34(2)**, 172-180.
- 117 Spinelli, S., Chefer, S., Suomi, S. J., Higley, J. D., Barr, C. S., & Stein, E. (2009). Early-life stress induces long-term morphologic changes in primate brain. **Archives of General Psychiatry**, **66(6)**, 658-665.
- 118 Barr, C. S., Newman, T. K., Shannon, C., Parker, C., Dvoskin, R. L., Becker, M. L., ... & Higley, J. D. (2004). Rearing condition and rh5-HTTLPR interact to influence limbic-hypothalamic-pituitary-adrenal axis response to stress in infant macaques. **Biological Psychiatry**, **55(7)**, 733-738.
- 119 Barr, C. S., Newman, T. K., Lindell, S., Shannon, C., Champoux, M., Lesch, K. P., ... & Higley, J. D. (2004). Interaction between serotonin transporter gene variation and rearing condition in alcohol preference and consumption in female primates. **Archives of General Psychiatry**, **61(11)**, 1146-1152.
- 120 Schwandt, M. L., Lindell, S. G., Higley, J. D., Suomi, S. J., Heilig, M., & Barr, C. S. (2011). OPRM1 gene variation influences hypothalamic-pituitary-adrenal axis function in response to a variety of stressors in rhesus macaques. **Psychoneuroendocrinology**, **36(9)**, 1303-1311.
- 121 Wendler, D. (2014). Should protections for research with humans who cannot consent apply to research with nonhuman primates?. **Theoretical Medicine and Bioethics**, **35(2)**, 157-173.